


# Neutrino Astronomy




This presentation is based on the excellent review of neutrino astronomy by Halzen & Klein 2010, and from one of his Plenary presentations in 2015.

# Neutrino Astronomy


- Neutrinos are ideal astronomical messengers. They travel from the edge of our observable Universe with little absorption and since they have no charge they are not deflected by magnetic fields.
- Unlike photons that are easily absorbed, high-energy neutrinos may reach us unscathed from cosmic distances, from the inner neighborhood of black holes.
- Because neutrinos are not easily absorbed they are very difficult to detect and large detector volumes of the order of a cubic-kilometer are required.

# Photon Spectrum of the Universe



# Neutrino Spectrum

- Solar neutrinos have energies of about 0.1 MeV to 10 MeV
- SN 1987A neutrinos (from the LMC) had energies  $\sim 100$  GeV
- AGN and GRB neutrino flux is higher than the “background” for neutrinos with energies above 100 TeV



# GZK Neutrinos


- Above a threshold of  $4 \times 10^{19}$  eV (1 TeV =  $10^{12}$  eV, 1 PeV =  $10^{15}$  eV ) cosmic rays (energetic charged particles) interact with the cosmic microwave background photons introducing an absorption feature in the cosmic-ray flux, the Greissen–Zatsepin–Kuzmin cutoff.
- Because of the GZK effect **cosmic rays with  $E > 4 \times 10^{19}$  eV can only travel for about 75 Mpc before they get absorbed** in the interaction:

COSMIC ray + CMB photon  $\rightarrow$  pion + neutron

- The charged pion will eventually decay into a GZK neutrino with an energy of  $> 10^6$  TeV
- The prediction is that the detection rate of GZK neutrinos is one per cubic kilometer per year. The direction of these neutrinos point back to the location of the source.



# Cosmic Ray Spectrum

- Cosmic accelerators can produce particles with energies in excess of  $10^8$  TeV
- The cosmic ray energy spectrum follows a sequence of three power laws.
- The first two are separated by a feature dubbed the “knee” at an energy of approximately 3000 TeV.
- There is evidence that cosmic rays up to this energy are galactic in origin.
- Cosmic rays with energies near and above the "ankle" are thought to be extragalactic in origin.



# Cosmic Ray Accelerators

- To produce cosmic rays requires protons to be accelerated to  $> \text{TeV}$  energies
- Cosmic Ray accelerators include Black Holes, Gamma Ray Bursts and Supernova Remnants
- The accelerated protons may interact with a photon to produce charged pions that latter decay into cosmic rays + neutrinos



# IceCube




ultra-transparent ice below 1.5 km

# IceCube


- The few neutrinos that interact with a nucleus in the ice create muons as well as electromagnetic and hadronic secondary particle showers.
- The charged particles that are produced from the interaction of the neutrino with the ice produce **Cherenkov Light** that spreads through the transparent ice.
- **Cherenkov light** is radiated by charged particles moving faster than the speed of light in the medium; in ice, this is 75% of the speed of light in a vacuum.
- **The Cherenkov light** is captured by photomultipliers (that use the photoelectric effect).
- The absorption length of light in regular tap water is about 1 m. The absorption length of light in the south pole ice is  $> 100\text{m}$ .
- The direction of the incoming neutrino can be recovered from the track left in the detector array.
- All detected events in the photomultiplier tubes are time-stamped to  $\sim 2$  nanoseconds precision.

# IceCube

## IceCube




# IceCube




muon track: color is time; number of photons is energy

# IceCube



# IceCube

- shielded and optically transparent medium
- muon travels from 50 m to 50 km through the water at the speed of light emitting blue light along its track



muon

interaction

• lattice of photomultipliers

neutrino


# IceCube

“Noise sources”: Atmospheric Neutrinos and Atmospheric Muons

- IceCube detects about 100 atmospheric neutrinos per day with energies above 100 GeV.
- Cosmic rays that collide with nitrogen and oxygen in our atmosphere create pions and kaons. The decay of charged pions and kaons produces the atmospheric neutrinos.
- Cosmic ray showers above IceCube will generate atmospheric muons and atmospheric neutrinos
- Cosmic ray showers in the atmosphere distant from IceCube will produce atmospheric neutrinos in IceCube (the muons decay before they get to IceCube into  $e^-$  and 2 neutrinos of different type)

# IceCube

## Signals and Backgrounds



# IceCube

muons detected per year:


- atmospheric\*  $\mu$   $\sim 10^{11}$
- atmospheric\*\*  $\nu \rightarrow \mu$   $\sim 10^5$
- cosmic  $\nu \rightarrow \mu$   $\sim 10$

\* 3000 per second

\*\* 1 every 6 minutes

# Discovery of Cosmic Neutrinos

cosmic neutrinos in 2 years of data at 3.7 sigma



# GZK neutrino search

GZK neutrinos are produced from the following process:

A cosmic ray ( $p$ ) with energy above  $\sim 4 \times 10^{19}$  eV interacts with a CMB photon ( $\gamma$ ) producing a pion ( $\pi$ ). The charged pion will decay into a neutrino ( $\nu$ ) and the muon will decay into a positron ( $e^+$ ) and neutrinos with  $\sim 10^6$  TeV energy!


$$p + \gamma \rightarrow n + \pi^+ \quad \text{and} \quad p + \pi^0$$

$$\pi^+ \rightarrow \mu^+ + \nu_\mu \rightarrow \left\{ e^+ + \nu_e + \bar{\nu}_\mu \right\} + \nu_\mu$$

# GZK neutrino search

GZK neutrino search: two neutrinos with  $> 1,000$  TeV


date: **August 9, 2011**  
energy: **1.04 PeV**  
topology: **shower**  
nickname: **Bert**



# Origin of Astrophysical Neutrinos

4 year HESE

**ICECUBE PRELIMINARY**



where do they come from?

# Origin of Astrophysical Neutrinos

- The 4 year IceCube results confirm the presence of astrophysical neutrinos and provide a measurement of the flux of astrophysical neutrinos.
- The sources of these neutrinos are expected to be black holes and massive exploding stars (GRBs). These sources are able to accelerate protons to  $> 10^6$  TeV
- The map of the detected locations of the astrophysical neutrinos indicates that most are of extragalactic origin but a small fraction may be from our own galaxy.
- Neutrinos above 100 TeV cannot be produced in the Earth's atmosphere indicating that these are astrophysical in origin.
- The neutrino-induced tracks are used to locate the position of the sources to an accuracy of less than a degree in the sky.

Where are the gamma rays that should accompany  
neutrinos?

$$p + \gamma \rightarrow n + \pi^+ \text{ and } p + \pi^0$$

$$\pi^+ \rightarrow \mu^+ + \nu_\mu \rightarrow \left\{ e^+ + \nu_e + \bar{\nu}_\mu \right\} + \nu_\mu$$

(charged pion decays  $\rightarrow$  cosmic rays and neutrinos)


$$\pi^0 \rightarrow e^- + e^+ + \gamma$$

(neutral pion decays  $\rightarrow$  cosmic rays and gamma rays)

# Where are the gamma rays that accompany neutrinos?

hadronic gamma rays ?

$$\pi^+ = \pi^- = \pi^0$$




$\gamma$


$\nu$

hadronic  
gamma rays


# Where are the gamma rays that accompany neutrinos?



# Where are the gamma rays that should accompany neutrinos? Answer: Fermi detects them





# Where are the gamma rays that should accompany neutrinos? Produced mostly by Blazars above 50 GeV



Global view of the diffuse  $\gamma$ -ray predictions is displayed for unresolved **BL Lacs** (**dotted green**) and for the sum of misaligned AGNs, star-forming galaxies, FSRQs, and millisecond pulsars (orange dashed line and uncertainty band). IGRB data are also displayed with black points. The sum of all the predictions is displayed in a blue curve line and cyan uncertainty band (figure from Di Mauro et al. 2014)

# number of muon neutrino events from gamma ray sources in 5 years





*This image shows one of the highest-energy neutrino events of this study superimposed on a view of the IceCube Lab (ICL) at the South Pole. Image: IceCube Collaboration*